

Anlage 3 Application of autologous dendritic cells in small animals according to the delimitation of responsibilities agreement

Name pet owner			
Name patient		Diagnosis	

1. Application using the Luer-Lock-System

For the application of cells, follow these steps:

- Mix gently by rotating the syringe.
- Screw off the cap.
- Attach the cannula on to the Luer-Lock.
- Apply the cells as described in the application instructions.

2. Intradermal Application Instructions

Intradermal application is the most common form of administering the dendritic cells. The application is often done in the inguinal area as this area has very few hair. Other locations for the application are also possible. Please remember to disinfect the location after shearing.

	
1. The cell suspension is filled in a sealed syringe.	2. Carefully swirl the cell suspension, twist off the cap and attach the cannula.
<p>Cleaning & Disinfection</p>	
3. Thoroughly clean and disinfect hands and injection site.	4. The number of wheals injected can be determined individually.

3. Confirmation

I hereby confirm that

- the autologous dendritic cell suspension was injected intradermally under sterile conditions as described above*, and
- I agree with the delimitation of responsibility for the implementation of immunotherapy with autologous dendritic cell suspension.

Date _____ Signature _____

*If individual work steps cannot be adhered to, please contact our team directly by phone +49-5522-9182581.

Please send this confirmation as an attachment by email to info@petbiocell.de

SOP TR-002/V-09	Anlage 3	Gültig ab: 30.06.2025	Seite: 1 von 1
Probenausgang			